

Content available at: <https://www.ipinnovative.com/open-access-journals>

The Journal of Community Health Management

Journal homepage: <https://www.jchm.in/>

Case Report

Pelvic floor physiotherapy for tenesmus-A case report

Binal Dave^{1*}, Rozina Shaikh¹

¹Dept. of Women's Health, Aqua Centric Private Limited, Mumbai, Maharashtra, India



ARTICLE INFO

Article history:

Received 29-09-2023

Accepted 26-10-2023

Available online 09-01-2024

Keywords:

Tenesmus

Pelvic floor dysfunction

Pelvic floor physiotherapy

Pelvic floor rehabilitation

ABSTRACT

Background: Tenesmus is uncomfortable sensation of incomplete evacuation of the bowel movements and it may be a distressing symptom for sufferers who suffer from rectal fullness. Not like nociceptive ache, neuropathic pain or visceral ache, little is thought about the pathway for this ache. There are only a few studies analyzing the palliative care of this symptom.

Case Presentation: In this case report, we describe a patient who suffered from severe tenesmus.

Case Management: To address the patient's tenesmus, pelvic floor physiotherapy was given.

Case Outcome: After 3 months of pelvic floor physiotherapy, the patient reported complete abolishment of her tenesmus symptoms.

Conclusion: Pelvic floor physiotherapy may be an effective tool for the management of tenesmus. Future research studies should consider examining the signs and symptoms of tenesmus.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Tenesmus is uncomfortable sensation of incomplete evacuation of the bowel movements¹ and it may be a distressing symptom for patients who be afflicted by rectal fullness.² The exact occurrence of this signs and symptoms in patients with rectal fullness is unclear, but based totally on our review, the superiority of intense tenesmus is quite low. Possibly due to this, tenesmus is an understudied symptom.³ In this case report, the authors described a new tool-pelvic floor physiotherapy for rectal fullness, rectal pain and incomplete evacuation and noted that the sensation of tenesmus additionally disappeared in all of the instances.⁴

2. Case Report

Our patient is a 27 year old girl with records of incomplete evacuation of bowel movements, stomach heaviness, rectal ache and rectal fullness since 6 months. Although she had

belly ache, she complained of severe tenesmus which she defined as an extreme and unrelenting choice to defecate.

For the control of her tenesmus, she tried on a variety of medicines such as clonazepam, ven-lafaxine, gabapentin, rectal diltiazem in addition to excessive dose opioids (oral hydromorphone 23 mg/day and oral methadone 10 mg/day) with marginal relief. For definitive control, she turned into trial of gemcitabine and carboplatin for which she became a non-responder and remedy was stopped. She was also considered for surgical therapy.

2.1. Case management

Appreciating the dearth of other alternative therapies, we carried out pelvic floor physiotherapy. Pelvic floor physiotherapy is a term which incorporates many different therapeutic approaches, together with however not constrained to electromyographic (EMG) biofeedback-guided pelvic floor muscle training (PFMT), that is the most extensively used as a rehabilitative treatment modality.

* Corresponding author.

E-mail address: doctorbinal85@gmail.com (B. Dave).

Pelvic floor physiotherapy is commonly administered by a pelvic floor physiotherapist, even though nurses, physicians, and other workforce can get hold of schooling to perform a lot of these interventions. The special rehabilitation techniques can be used independently, However in conjunction with one another as a holistic method to supply the maximum gain for the affected person.⁵ The primary aim of all types of pelvic floor rehabilitation is to improve pelvic floor and anal sphincter muscle strength, tone, endurance, and coordination for reduction in signs and symptoms. Additional benefit is that a person's recognition of their own pelvic floor muscle groups, enhancing rectal sensitivity, rectal fullness, incompleteness of evacuating stools and decreasing scar burden to allow for advanced muscle characteristic.

Pelvic floor physiotherapy techniques encompass bowel management education and management retraining, PFMT, biofeedback therapy (BFT), using electrical stimulation guided myofascial release and connective tissue mobilization strategies.^{6–9}

Bowel education and retraining can include many one-of-a-kind factors, a focal point on life-style changes along with instruction as to take greatest fluid intake and nutritional modifications may be important for a affected person. PFMT describes any quantity of various processes for growing strength, endurance, and coordination of the pelvic floor muscles and anal sphincters. Thoracoabdominopelvic muscle education has additionally been endorsed, as it has been theorized that education of all center muscle tissues to work in tandem might be extra effective than a slender cognizance on the pelvic ground muscle groups alone. A unique interest is regularly paid to the transversus abdominus in such improved strategies. PFMT commonly includes verbally guided coaching in pelvic floor and sphincter contractions (Kegel contractions). Sufferers can be taught to settlement in a ramification of approaches. A few examples consist of maximal voluntary sustained sphincter contractions, submaximal sustained contractions, and rapid-twitch or quick-flick contractions. A commonly pronounced PFMT approach is to examine the pelvic floor to an elevator, capable of stop at exclusive floors as it ascends and descends. Different strategies encompass working on coordination of anal sphincter activity and working to isolate a contraction of the anal sphincter. A few practitioners use their hand located externally, or a digit located vaginally or rectally to help educate the affected person in the best exercising techniques, but most would argue that this constitutes a form of low-tech biofeedback schooling.^{10–13}

2.2. Case outcome

In the therapy room, the patient was assessed and mentioned to have complete relief of her tenesmus. She scored the severity of the tenesmus as 0/10. We stated that at the 1-

week and 3 monthly follow up, there was no recurrence of her tenesmus symptoms. however, she persisted to consume the identical amount of opioids as her belly ache have become greater prominent after the tenesmus signs and symptoms had resolved. On an average, she said that she become relatively satisfied with the pelvic floor physiotherapy There have been no quick-time period or long-term damaging effects.

3. Conclusion

Pelvic rehabilitation methods including PFMT, biofeedback guided pelvic floor relaxation, biofeedback guided rectal sensitivity and coordination education, and electric stimulation may be effective tools in the management of tenesmus Further research is needed to outline the function of rehabilitation, predictors of desirable results, and the most efficacious remedy protocols.

4. Source of Funding

None.

5. Conflict of Interest

None.


References

1. Bols EM, Berghmans BC, Hendriks EJ. A randomized physiotherapy trial in patients with fecal incontinence: design of the PhysioFIT-study. *BMC Public Health*. 2007;7:355. doi:10.1186/1471-2458-7-355.
2. Perry S, Shaw C, Mcgrother C. Prevalence of faecal incontinence in adults aged 40 years or more living in the community. *Gut*. 2002;50(4):480–4.
3. Tan JJ, Chan M, Tjandra J. Evolving therapy for fecal incontinence. *Dis Colon Rectum*. 2007;50(11):1950–67.
4. Van Oud-Alblas M, Thomeer BJ, Stam HJ, Van Overbeeke A, Consten E. Fecal incontinence: an update on available techniques in diagnosis and treatment. *Surg Technol Int*. 2008;17:156–64.
5. Norton C, Whitehead WE, Bliss DZ, Harari D, Lang J. Management of fecal incontinence in adults. *Neurolog Urodyn*. 2010;29(1):199–206.
6. Van Tets W, Kuijpers JH, Bleijenberg G. Biofeedback treatment is ineffective in neurogenic fecal incontinence. *Dis Colon Rectum*. 1996;39(9):992–4.
7. Norton C, Cody J. Biofeedback and/or sphincter exercises for the treatment of faecal incontinence in adults. *Cochrane Datab Syst Rev*. 2012;11(7):CD002111. doi:10.1002/14651858.CD002111.pub3.
8. Hosker G, Norton C, Brazzelli M. Electrical stimulation for faecal incontinence in adults. *Cochrane Database Syst Rev*. 2000;2:CD001310.
9. Santoro GA, Wieczorek AP, Bartram CI. Imaging and Multidisciplinary Approach to Management. In: and others, editor. *Pelvic Floor Disorders*. Springer-Verlag Italia; 2010. p. 317.
10. Norton C, Chelvanayagam S. Methodology of biofeedback for adults with fecal incontinence: a program of care. *J Wound Ostomy Continence Nurs*. 2001;28(3):156–68.
11. Hansen JL, Bliss DZ, Mcalpine CP. Diet strategies used by women to manage fecal incontinence. *J Wound Ostomy Continence Nurs*. 2006;33(1):61–2.
12. Bliss DZ, Jung HJ, Savik K. Supplementation with dietary fiber improves fecal incontinence. *Nurs Res*. 2001;50(4):203–13.

13. Stokes G. Bowel Care in Older People. and others, editor. Clinical Effectiveness & Evaluation Unit, Royal College of Physicians; 2002. p. 97–109.

Rozina Shaikh, Senior Physiotherapist

Author biography

Binal Dave, Professor  <https://orcid.org/0000-0001-8512-5627>

Cite this article: Dave B, Shaikh R. Pelvic floor physiotherapy for tenesmus-A case report. *J Community Health Manag* 2023;10(4):163-165.