

AN IMMUNOLOGICAL STUDY OF BRONCHIAL ASTHMA WITH SPECIAL REFERENCE TO INTERLEUKIN-4, INTERFERON- γ AND IMMUNOGLOBULIN E

J K Mishra,¹ Rashid Parvez,² Usha,³ Girish Singh⁴

1. Professor, Department of T.B. & Respiratory Diseases, IMS, Banaras Hindu University, Varanasi.
2. Department of T.B. & Respiratory Diseases, IMS, Banaras Hindu University, Varanasi.
3. Professor, Department of Pathology, IMS, Banaras Hindu University, Varanasi.
4. Division of Biostatistics, Institute of Medical Sciences, Banaras Hindu University, Varanasi, India

Corresponding Author:

E-mail: jkmishra.imsbhu@gmail.com

ABSTRACT

Asthma is a disorder defined by its clinical, physiological, and pathological characteristics. The predominant feature of the clinical history is episodic shortness of breath, particularly at night, often accompanied by cough. The main physiological feature of asthma is episodic airway obstruction characterized by expiratory airflow limitation. The dominant pathological feature is airway inflammation, sometimes associated with airway structural changes. This immunological study on bronchial asthma was conducted at Sir Sunderlal Hospital, Institute of Medical Sciences, Banaras Hindu University, Varanasi in the year 2010. Based on clinical history one group of 70 cases with history of asthma and other group of 36 controls with no history of asthma were made. The mean serum level of IL-4, IgE, and AEC was observed significantly higher in cases as compared to controls whereas; mean IFN- γ as well as FEV₁ were observed significantly lower in cases as compared to controls. The cut-off values at appropriate sensitivity and specificity using ROC curve for IL-4, IFN- γ , IgE, AEC and FEV₁ were found as 7.25, 0.35, 225, 345 and 79.50 respectively. The imbalance in the serum levels of IL-4 and IFN- γ is predicted to drive the asthma pathogenesis while reducing IgE levels may now be considered as a treatment strategy.

Key-words: IL-4, IFN- γ , serum IgE, bronchial hyper-responsiveness, AEC, FEV₁

Introduction

Asthma is a common, chronic disease of airway inflammation that manifests with recurrent episodes of coughing, breathlessness, wheezing, and chest tightness. These episodes are associated with airflow obstruction that is at least partially reversible. Making the diagnosis of asthma requires a clinical evaluation of the patient's symptom, medical history, physical examination and diagnostic tests.

Asthma is a problem worldwide, with an estimated 300 million affected individuals. Despite hundreds of reports on the prevalence of asthma in widely differing populations, the lack of a precise and universally accepted definition of asthma makes reliable comparison of reported prevalence from different parts of the world problematic.

Nonetheless, based on the application of standardized methods to measure the prevalence of asthma and wheezing illness in children and adults it appears that the global prevalence of asthma ranges from 1% to 18% of the population in different countries. There is limited data on asthma epidemiology from the developing world, including India. Although some attempts have been made, studies suffer from several scientific drawbacks including lack of uniformity of methodology and analysis of data. Asthma rates are officially low in India, although there is some recent evidence that the true prevalence is higher than previously thought. To date, the total estimated burden of Asthma is an overall prevalence of 3% (30 million patients), and among adults over the age of 15, a median prevalence of 2.4%. [1]

The factors influencing the development and expression of asthma may be classified into two categories viz., host factors (genetic, obesity, sex) and environmental factors (Allergens, Infections, Occupational sensitizers, Tobacco smoke, Diet). Asthma being an inflammatory disorder of the airways, involves several inflammatory cells and multiple mediators that result in characteristic pathophysiological changes. In ways that are still not well understood, this pattern of inflammation is strongly associated with airway hyper-responsiveness and asthma symptoms. Over 100 different mediators are now recognized to be involved in asthma and mediate the complex inflammatory response in the airways. [2] The mucosal mast cells activated by allergens through high-affinity Immunoglobulin E (IgE) receptors, as well as by osmotic stimuli resulting to increased in numbers in airway smooth muscle may be linked to airway hyper-responsiveness. [3] Eosinophils and T lymphocytes present in increased numbers in the airways, release specific cytokines, including Interleukin-4 (IL-4) that orchestrate eosinophilic inflammation and IgE production by B lymphocytes. [4,5] Interaction of dendritic cells with regulatory T cells ultimately stimulate production of Th-2 cells from naive T cells. [6,7]

Thus determination of serum levels of IL-4 and IFN- γ may be useful for understanding and monitoring the inflammatory response in asthma. Keeping in view of these facts the aim of the present work was to evaluate serum level of Interleukin-4, Interferon gamma and Immunoglobulin E in Bronchial asthma and study their interrelationship. Further, also to determine

the cutoff value of IL-4, IFN- γ and IgE, through ROC curve and find sensitivity and specificity for the chosen cutoff.

Material and Method

The present study based on total 106 subjects was conducted at Sir Sunderlal Hospital, Institute of Medical Sciences, Banaras Hindu University, Varanasi in the year 2010. Based on the clinical history of investigations two group were made. Group-1 consisted of 70 cases who had history of asthma (history of allergy and spirometry with post bronchodilation reversibility $\geq 12\%$ and $\geq 200\text{ml}$ (FEV1). Group-2 had 36 controls that had no history of asthma.

Patient selection

Inclusion Criteria:

1. Cases of Br. Asthma-Cases taken fulfill the criteria of bronchial Asthma (h/o allergy and spirometry with post- bronchodilation reversibility $\geq 12\%$ and $\geq 200\text{ml}$ (FEV1).

Exclusion Criteria:

1. No h/o Br. Asthma, allergy, viral infection, HIV, alcohol intake parasitic, infestation severe burn cases Churg-Strauss syndrome and Wiskott -Aldrich syndrome.
2. Any other respiratory co-morbidity like Pneumonia, Pul. Koch's.
3. Any other systemic co-morbidity like hypertension, diabetes.
4. Patient not giving consent.

The diagnosis and clinical evaluation

A detailed medical history of patient known or thought to have asthma was taken. Present history, history of past illness, family history, and treatment history were recorded. Clinical examinations done including general examination, systemic examination, URT, chest, auscultation, skin examination, CVS examination, CNS examination, routine haematological investigation and X-ray chest PA view.

The diagnostic test for Absolute Eosinophil Count, pulse oxymetry, serum level of IgE, ABG (Arterial blood gas analysis), sputum for fungal mycelia, stool examination for parasites, Pulmonary Function Test (PFT) with reversibility $\geq 12\%$ and $\geq 200\text{ ml}$ from pre-bronchodilator value were performed. Serum was separated from the blood, which was aseptically drawn by venipuncture, after clotting and centrifugation the sample immediately transferred in plastic tube and stored at $< -18^{\circ}\text{C}$ maximum for 2 month or preferably at $< -70^{\circ}\text{C}$ for longer time. Serum IL-4, serum IFN- γ and serum IgE were estimated by ELISA technique. These investigations were done in Immuno-Pathology Department, IMS-BHU. Absolute Eosinophil Count (AEC) was done in CCI lab, S.S. Hospital BHU. Pulmonary Function Test (PFT) was done in TB and Respiratory Department with Medisoft PFT machine.

Statistical Analysis

Analysis of data was done using SPSS ver.16 statistical software package. Cross tables, mean and standard deviation (sd) for both groups were obtained and inter group comparison was done. ROC curve with area under the curve was used to determine the cutoff value of IL-4, IFN- γ and IgE. Sensitivity and specificity for the chosen cutoff were also determined. To determine the degree of linear interrelationship between these variables, coefficient of correlation was also determined.

Result

The present study was based on total 106 subjects. The number of cases in Group-1 (cases with history of Bronchial Asthma) was 70 whereas; in Group-2 (control group with no history of Bronchial Asthma) the number was 36. The percentage of males in Group-1 was 54.3 while of females it was 45.7. Similarly the percentage of males and females in Group-2 was 63.9 and 36.1, respectively (**Table-1**). In this study age range for asthmatic cases was 10-70 yrs (mean age 30.78 ± 11.27) and for control group it was 10-60 yrs (mean age 29.23 ± 12.94). The difference between mean ages of the two groups was statistically not significant (**Table-2**).

Table-1: Age and sex distribution according to groups

Variables		Group-1	Group-2
		Case	Control
Number		70	36
Sex	Male	38 (54.3%)	23 (63.9%)
	Female	32 (45.7%)	13 (36.1%)
Age (yrs)	10 - 20	07	10
	21 - 30	33	12
	31 - 40	21	07
	41 - 50	04	05
	51 - 60	04	02
	61 - 70	01	00

The mean IL4 value for cases (10.16 ± 2.69) was found higher than those of control group (8.06 ± 4.01) and this difference between means was statistically significant ($p < 0.01$). Similarly mean IgE value for cases (236.70 ± 73.21) was found higher than those of control group (209.33 ± 42.60) and this difference was statistically significant ($p < 0.05$). The mean value of AEC for case group was $464.31 (\pm 121.58)$ while it was $223.53 (\pm 83.37)$ for control group and the inter group comparison resulted statistically significant ($p < 0.001$). The mean IFN- γ for cases (0.390 ± 0.266) was observed significantly lower ($p < 0.02$) than that of control group (0.518 ± 0.314). Similarly, mean FEV₁ value for cases (75.63 ± 4.68) was found lower than that of control group (82.36 ± 1.92) and this difference was statistically significant ($p < 0.001$) (**Table-2**).

Table-2: Mean \pm SD and Statistical Significance

Variables	Group-1: Case	Group-2: Control	*Between the group comparison (p-value)
	Mean \pm SD	Mean \pm SD	
Age (yrs.)	30.78 \pm 11.27	29.23 \pm 12.94	0.471 (N.S.)
Interleukin-4(pg/ml)	10.16 \pm 2.69	8.06 \pm 4.01	0.001
IFN- γ (iu/ml)	0.39 \pm 0.266	0.518 \pm 0.314	0.011#
IgE(iu/ml)	236.70 \pm 73.21	209.33 \pm 42.60	0.041
AEC(cells/cumm)	464.31 \pm 121.58	223.53 \pm 83.37	0.000
FEV ₁	75.63 \pm 4.68	82.36 \pm 1.92	0.000

* Independent Sample t-test, # Wilcoxon-Mann-Whitney test

To measure the degree of linear relationship between IL-4, IFN- γ , IgE, FEV₁ and AEC Karl Pearson coefficient of correlation and its statistical significance was also determined. A negative correlation coefficient though, not statistically significant ($p > 0.05$) was observed between IL4 and IgE, IL4 and AEC, IFN- γ and IgE, IFN- γ and AEC, IgE and FEV1. Whereas, a positive correlation coefficient though, not statistically significant ($p > 0.05$) was observed between IL4 and IFN- γ , IL4 and FEV1, IgE and AEC. It is to be noted that correlation coefficient between IFN- γ and FEV1 was positive and statistically significant ($p < 0.05$) whereas, correlation coefficient between AEC and FEV1 was negative and statistically significant ($p < 0.05$).

Table-3: ROC area under the curve, cut-off value, sensitivity and specificity

Variable	ROC Area under the curve	Std. Error	Cut-off value	Sensitivity	Specificity
IL-4 (pg/ml)	0.709	0.061	7.25	84.3%	58.3%
IFN- γ (iu/ml)	0.644	0.060	0.350	74.3%	64.1%
IgE (IU/ml)	0.676	0.055	225	62.9%	66.7%
AEC (Cells/cumm)	0.958	0.016	345	84.3%	97.2%
FEV ₁ (%)	0.962	0.022	79.50	95.7%	100%

ROC curve was plotted and area under the curve, cut-off value of the tests and its sensitivity, specificity at that cut-off were determined and shown in **Table-3**. The cut-off values for IL-4, IFN- γ , IgE, AEC and FEV₁ were 7.25, 0.35, 225, 345 and 79.50 respectively.

Discussion

Asthma is a chronic inflammatory disorder of the airways in which many cells and cellular elements play a role. The chronic inflammation is associated with airway hyper-responsiveness that leads to recurrent episodes of wheezing, breathlessness, chest tightness, and coughing, particularly at night or in the early morning. These episodes are usually associated with widespread, but variable, airflow obstruction within the lung that is often reversible either spontaneously or with treatment. Recent investigations have demonstrated that allergen-specific CD4+T helper 2 (Th-2)-type lymphocytes and the cytokine interleukin (IL)-4 play important roles in the development of airway eosinophilic inflammation and bronchial hyper-responsiveness (BHR). Interleukin-4 is thought to be involved in allergic reactions, because it induces the differentiation of naive CD4+ T lymphocytes into Th-2 type cells,[8] mast cell proliferation[9] and also IgE synthesis.[10] The present study showed increased mean serum level of IL-4 for cases of bronchial asthma in comparison to healthy control. A number of authors have studied serum cytokines in asthmatic subjects. Bogic´ et al[11] have

reported significant higher IL-4 serum concentrations in asthmatic group compared to control and these were significantly higher in patients with moderate and severe asthma compared to mild asthmatics. Shahid et al[12] had shown an increased concentration of exhaled IL-4 in asthmatic and decreased IFN- γ . However, results from recent studies regarding the role of IL-4 in the onset of allergen-induced BHR are still conflicting. [13]

In the present study the mean serum level of IFN- γ was observed significantly lower as compared to control group. It had been demonstrated in this study that patient with bronchial asthma (atopic) produce significantly less IFN- γ and more IL-4. IFN- γ has been responsible for regulating the activation, differentiation and recruitment of eosinophil.

In allergic bronchial asthma, IgE is thought to play an important role in the generation of mediators and in the development of BHR.[14] Brusselle et.al[15] reported that aeroallergen-induced airway inflammation and BHR are dependent on IL-4 in gene disrupted mice. Several other studies have also reported that allergic and asthmatic subjects are more likely to have elevated levels of Th-2 cytokines and reduced levels of Th-1 cytokines. It could

be argued that Th-1/Th-2 imbalance is responsible for the development of IgE-mediated inflammation. The present study also revealed significantly increased mean serum levels of IgE in comparison to control group. The serum IgE level was observed negatively correlated with the amount of IFN- γ . Further, the mean Absolute Eosinophil Count (AEC) of cases with bronchial asthma was found significantly higher in comparison to control group. The reason of observing significantly lower mean FEV₁ in cases of Bronchial Asthma as compared to control group may be due to increased BHR and obstruction in cases. The association between eosinophils and allergic disease has been known for many years. According to Halonen *et.al*, [16] a significant relationship exists between serum IgE levels and eosinophilia in populations presumed to be free of parasites where IgE levels presumably provide a better clue to atopy than do skin tests. This shows that increased bronchial hyper-responsiveness and obstruction occur if serum level of AEC and IgE are raised. Thus IgE secretion by lymphocytes defines the allergic state and nearly all asthmatics have a higher IgE levels in serum than normal, following adjustment with age and sex.

Conclusion

Bronchial asthma is characterized by lower respiratory tract inflammation leading to

bronchial hyper-responsiveness (BHR) with variable and reversible airflow obstruction. Asthma has significant genetic and environmental components, but since its pathogenesis is not clear, much of its definition is descriptive. Interleukin-4 being the main cytokine involved in the pathogenesis of allergic responses that can also down-regulate acute inflammatory changes, also got additional effects on asthma pathogenesis which include stimulation of mucus producing cells and fibroblasts leading to airway remodeling. It has also been confirmed that the crucial role of IL-4 lies in its effect on Th-2 development, rather than on the induction of IgE synthesis and subsequent mast cell degranulation. On the other hand IFN- γ is a potent inhibitor of IgE synthesis. Thus, this imbalance in the serum levels of IL-4 and IFN- γ is predicted to drive the asthma pathogenesis. The findings of present study support the hypothesis of Th-1/Th-2 cytokine imbalance and suggest that serum level of IL-4 may be elevated in concert with decreased level of IFN- γ in asthma. Determination of serum levels of IL-4 and IFN- γ may be useful for understanding and monitoring the inflammatory response in asthma. Reducing IgE levels may now be considered as a strategy for the treatment of allergic rhinitis and asthma both.

References

1. Aggarwal AN, Chaudhry K, Chhabra SK, et al. Prevalence and risk factors for bronchial asthma in Indian adults: a multicentre study. *Indian J Chest Dis Allied Sci.* 2006; 48:3-22
2. Galli SJ, Kalesnikoff J, Grimbaldston MA, Piliponsky AM, Williams CM, Tsai M. Mast cells as "tunable" effector and immunoregulatory cells: recent advances. *Annu Rev Immunol* 2005; 23:749-86.
3. Robinson DS. The role of the mast cell in asthma: induction of airway hyper-responsiveness by interaction with smooth muscle? *J Allergy Clin Immunol.* 2004; 114 (1): 58-65.
4. Kay AB, Phipps S, Robinson DS. A role for eosinophils in airway remodelling in asthma. *Trends Immunol* 2004; 25(9): 477-82.
5. Larche M, Robinson DS, Kay AB. The role of T lymphocytes in the pathogenesis of asthma. *J Allergy Clin Immunol* 2003; 111 (3): 450-63.
6. Kuipers H, Lambrecht BN. The interplay of dendritic cells, Th2 cells and regulatory T cells in asthma. *Curr Opin Immunol.* 2004; 16 (6) : 702-8.
7. Akbari O, Faul JL, Hoyte EG, Berry GJ, Wahlstrom J, Kronenberg M, *et al.* CD4+ invariant T-cell-receptor+ natural killer T cells in bronchial asthma. *N Engl J Med.* 2006; 354 (11): 1117-29.
8. Abehsira-Amar O, Gibert M, Jolij M, Theze J, Jankovic DG. IL-4 plays a dominant role in the differential development of Th0 into Th1 and Th2 cells. *J. Immunol.* 1992; 148: 3820-9.
9. Abehsira-Amar O, Gibert M, Jolij M, Theze J, Jankovic DG. IL-4 plays a dominant role in the differential development of Th0 into Th1 and Th2 cells. *J. Immunol.* 1992; 148: 3820-9.
10. Snapper CM, Paul WE. Interferon-g and B cell stimulatory factor-1 reciprocally regulate Ig isotype production. *Science* 1987; 236: 944-7.
11. Bogic´ M, Savic´ N, Jovic´ ic´ Z´ , Spiric´ VT, Popadic´ AP, Raškovic´ S, et al. Clinical significance of measurement of interleukin 4 and interleukin 5 serum concentrations in bronchial asthma. *Jugoslav Med Biochem* 2004; 23: 51-4.
12. Shahid SK, Kharitonov SA, Wilson NM, Bush A, Barnes PJ. Increased interleukin-4 and decreased interferon-c in exhaled breath condensate of children with asthma. *Am J Respir Crit Care Med* 2002; 165: 1290-3.
13. Hogan SP, Mould A, Kikutani H, Ramsay AJ, Foster PS. Aeroallergen-induced eosinophilic inflammation, lung damage, and airways hyper-reactivity in mice can occur independently of IL-4 and allergen-specific immunoglobulins. *J. Clin. Invest.* 1997; 99: 1329-39.
14. Mishra JK, Usha, Deb A, Sehgal D. A study of serum IgE in allergic diseases. *Indian J Allergy Asthma Immunol* 2009; 23(1): 29-36.
15. Brusselle G, Kips J, Joos G, Bluethman H, Pauwels R. Allergen-induced airway inflammation and bronchial responsiveness in wild type and interleukin-4 deficient mice. *Am. J. Respir. Cell Mol. Biol.* 1995; 12: 254-9.
16. Halonen M, Barbee RA, Lehanitz MD, Eurrows B. An epidemiologic study of the interrelationship of total serum immunoglobulin E, Allergy skin reactivity and eosinophils. *J Allergy Clin. Immunol.* 1982; 69: 221-228.

