

Need for public awareness: Peptic ulcer disease

J. Mukhopadhyay

Professor, Dept. of Community Medicine, JIMS Hospital, Budge Budge, Kolkata, West Bengal, India

*Corresponding Author: J. Mukhopadhyay

Email: jmukho@yahoo.co.in

Abstract

Peptic ulcer disease (PUD) which existed around the world has shown a decline both in developed and developing countries in the current decade due to improvement in sanitary conditions and availability of specific treatment along with proton pump inhibitors. PUD has been linked to *H. pylori* infection and rampant use of NSAIDs. Gastric Carcinoma is a fatal outcome of PUD which takes long time to develop providing a valuable window for prevention. Gastric Carcinoma is the fifth most common malignancy and third leading cause of cancer fatality. 80% of gastric carcinoma has been incriminated to *H. pylori* infection. Chronic *H. pylori* induced gastritis is preventable through observance of domestic, food and personal hygiene thereby averting faecal-oral and oral-oral transmission of the organism among family and community members. Informative awareness program supported with screening, surveillance, treatment and community outreach schedule has been recommended to reduce the morbidity and mortality burden of PUD.

Keywords: PUD, Gastric CA, *H. pylori*, PPI.

Peptic ulcer disease (PUD) has demonstrated a variable trend in the past. In the West, the incidence of duodenal ulcer intensified sharply at the turn of the 19th to 20th century but exhibited a perceptible decline later during the last three decades.^{1,2} In the East, the surge was equally notable to begin with, the decline appeared to have been delayed; started waning in the past decade only.^{3,4} The drift has been imputed to an enhancement in sanitary condition, better water supply and personal hygiene associated with widespread use of proton pump inhibitors (PPI) resulting in a declining trend of PUD.^{5,6} A global overview of 2018 recounted the highest annual incidence of all PUD as 141.8 per 100,000 population in Spain and the lowest 23.9 in UK. Highest annual incidence of bleeding PUD was 72.5 per 100,000 persons in Greece as compared to lowest of 8.3 in UK. Highest yearly incidence of perforated PUD has been 4.4 per 100,000 individuals in South Korea in contrast to lowest of 2.2 in UK.⁷ This indicates that the disease entity remains common around the world; however, existing regional data gap calls for further studies to grasp the insight.

Etiology of PUD includes *H. pylori* infection, NSAIDs, pepsin, smoking, alcohol, bile-acids, steroids, stress, changes in gastric mucin consistency and Crohn disease.^{8,9} However, the pathogenic role of *H. pylori* in chronic active gastritis and its association with duodenal ulcer in 95 to 99% of patients has been well established.¹⁰ Prevalence of *H. pylori* is higher in developing countries that estimated to have infected about 70% of the population, as compared to 40% in the developed countries.¹¹ In developing countries, where most children become infected by the age 10, gastric cancer rates are very high.¹² Despite use of eradication therapy for *H. pylori* infection along with PPIs and advanced endoscopic intervention during recent decades, mortality associated with PUD has not decreased concurrently with the fall in incidence.¹³ Worldwide, gastric cancer is the fifth most common cancer and third leading cause of cancer-related death.¹⁴ It has been estimated that

80% of all gastric cancer cases can be attributed to chronic *H. pylori* infection.¹⁵ The development of gastric cancer from *H. pylori*, which involves a multistep process from chronic to atrophic gastritis to mucosal metaplasia to dysplasia to gastric carcinoma, can take decades to develop.¹⁶ This slow progression provides an opportunity for early detection and treatment of *H. pylori* that may result in prevention of gastric cancer.

Risk factors of *H. pylori* infection in children are allied to ethnicities, geographic location, living conditions, personal hygiene, drinking water sources, type of housing, sewage system and garbage collection within the living environment. These risk factors are usually associated with socioeconomic status (SES) of the family.¹⁷ Specifically, lower SES, consumption of restaurant food, meat, fish, non-filtered water, smoking and family history of PUD are risk factors for *H. pylori* infection.¹⁸ Since mother-child and sib-sib transmission are most significant, the best way to avert *H. pylori* infection is to educate women about personal hygienic measures that can protect themselves and their offspring. Considering high morbidity and mortality load, significant infectivity, large population at risk, long disease progression period and simple preventive measures, it is felt pertinent that dissemination of awareness about PUD would benefit population at risk and prevent occurrence of gastric carcinoma at large. Awareness program needs to be buttressed by screening, surveillance, treatment and outreach schedule for the community.

Conflict of Interest: None.

References

1. Sonnenberg A. Temporal trends and geographical variations of peptic ulcer disease. *Aliment Pharmacol Ther* 1995;9:3-12
2. Baron JH, Sonnenberg A. Hospital admissions for peptic ulcer and indigestion in London and New York in the 19th and early 20th centuries. *Gut* 2002;50:568-70.
3. Goh KL, Wong HT, Lim CH, Rosaida MS. Time trends in peptic ulcer, erosive reflux esophagitis, gastric and

- oesophageal cancers in a multiracial Asian population. *Aliment Pharmacol Ther* 2009;29:774–80
4. Goh KL. Changing trends in gastrointestinal disease in the Asia-Pacific region. *J Dig Dis* 2007;8:179–85.
 5. Sonnenberg A. Causes underlying the birth-cohort phenomenon of peptic ulcer: analysis of mortality data 1911–2000, England and Wales. *Int J Epidemiol* 2006;35:1090-97.
 6. Pérez-Aisa MA, Del Pino D, Siles M, Lanás A. Clinical trends in ulcer diagnosis in a population with high prevalence of *Helicobacter pylori* infection. *Aliment Pharmacol Ther* 2005;21:65-72.
 7. H Azhari, F Underwood, J King, S Coward, S Shah, S Ng, G Ho, C Chan, W Tang, G G Kaplan. The global incidence of peptic ulcer disease and its complications at the turn of the 21st century: A systematic review. *J Can Assoc Gastroenterol* 2018;1(2):61–62. doi: <https://doi.org/10.1093/jcag/gwy009.036>
 8. Mynatt RP, Davis GA, Romanelli F. Peptic Ulcer Disease: Clinically Relevant Causes and Treatment. *Orthop* 2009;32(2):104
 9. Niv Y. H. pylori/NSAIDS - Negative Peptic Ulcer the Mucin Theory. *Med Hypotheses* 2010;75(5):433-5
 10. Wirth H, Yang M, Peek R, Tham K, Blaser M. *Helicobacter pylori* Lewis expression is related to the host Lewis phenotype. *Gastroenterol* 1997;113:1091-8
 11. Brown LM. *Helicobacter Pylori*: Epidemiology and Routes of Transmission. *Epidemiol Rev* 2000;22(2):283-97
 12. Telford J, Covacci A, Ghiara MP, Montecucco C, & Rappuoli R. Unravelling the pathogenic role of *Helicobacter pylori* in peptic ulcer: Potential new therapies and vaccines. *Trends Biotech* 1994;12(10):420-6.
 13. Malmi H. Peptic ulcer: incidence, associated morbidity and mortality (PhD). University of Helsinki, 2018 https://helda.helsinki.fi/bitstream/handle/10138/232277/PEPTI_CUL.pdf
 14. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global Cancer Statistics, 2012. *CA Cancer J Clin* 2015;65(2):87–108. doi:10.3322/caac.212623
 15. IARC *Helicobacter pylori* Working Group. *Helicobacter Pylori Eradication as a Strategy for Preventing Gastric Cancer*. Lyon: International Agency for Research on Cancer 2014. Available from: https://www.iarc.fr/wp-content/uploads/2018/07/pr227_E.pdf
 16. Correa P. Human gastric carcinogenesis: a multistep and multifactorial process – First American Cancer Society Award Lecture on Cancer Epidemiology and Prevention. *Cancer Res* 1992;52(24):6735–40. PMID: 1458460
 17. Ozbey G, Hanafiah A. Epidemiology, Diagnosis, and Risk Factors of *Helicobacter pylori* Infection in Children. *Eurasian J Hepatogastroenterol* 2017;7(1):34-9. doi: 10.5005/jp-journals-10018-1208
 18. Mhaskar RS, Ricardo I, Azliyati A, Laxminarayan R, Amol B, Santosh W, Boo K. Assessment of Risk Factors of *Helicobacter Pylori* Infection and Peptic Ulcer Disease. *J Glob Infect Dis* 2013;5(2):60–67. doi: 10.4103/0974-777X.112288

How to cite this article: Mukhopadhyay J. Need for public awareness: Peptic ulcer disease. *J Community Health Manag* 2019;6(2):64-5.