

The Journal of Community Health Management

Journal homepage: https://www.jchm.in/

Original Research Article Post-COVID impact: Misery of the indisposed

J Mukhopadhyay^{1,*}

¹Dept. of Community Medicine, Jagannath Gupta Institute of Medical Science & Hospital, Budge-Budge, Kolkata, West Bengal, India



PUBL

ARTICLE INFO

Article history: Received 05-04-2023 Accepted 19-04-2023 Available online 28-04-2023

Keywords: COVID -19 Medico-social problems Psychological assistance

ABSTRACT

Background: COVID-19 seized human life with sickness & fatality; even elicited medical, social and financial unrest. Clinical symptoms continued for weeks after contracting COVID-19 with socio-emotional turmoil. Considering these adversities, it was decided to ascertain medico-social effect of COVID-19 on patients coming in contact with scholar during May 2020 to September 2021 in Kolkata.

Materials and Methods: 83 COVID-19 positives were contacted for enlisting & consent on first opportunity, however only 79 responded & continued. A structured questionnaire was prepared including personal attributes, details of COVID-19 infection and post-COVID medico-social problems. Questionnaires were distributed online as and when contact established during study. Participants were contacted for clinical conditions every week during first month, thereafter once a month till 3 months. Finally, data collected were analyzed and tabulated.

Results: 34 & 45 subjects had COVID-19 during 2020 & 2021 respectively. Majority were educated employed males of 41-50 years with co-morbidities (35.6-52.9%). All followed COVID appropriate behaviour. Fever, fatigue, breathlessness, sore-throat & cough were common during first week; even continued for some in following weeks. Headache, anosmia & ageusia, giddiness, weight-gain & insomnia were also persistent. Isolation, anxiety, avoidance by relatives, hardship in daily needs with apathy of attendants made lingering problems. 13-35% had financial hang-ups due to expenses & helplessness. Time with family, online counsellor support and upgraded home delivery services with containment-respite were in high demand.

Conclusion: Study revealed prolonged medico-social sufferings of the COVID affected seeking online medical-social-psychological support. Further research is contemplated to plan possible future guidelines to resolve these kinds of hardships in pandemic.

This is an Open Access (OA) journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The catastrophic COVID-19 pandemic annihilated human life in an insurmountable manner in the recent past. The nefarious contagion denigrated human existence not only with disease & death but even triggered medical, social and economical disquiet among the afflicted. Patients continue to experience certain symptoms weeks and months after onset of COVID-19 regardless of disease severity along with much social & emotional quandary.^{1–3} Considering this disturbing tribulation, it was thought prudent to ascertain the detrimental effect of COVID-19 infection on medico-social front for the affected individuals those intersected with the scholar in personal and professional domain during May 20 to Sep 21 in Kolkata, India.

* Corresponding author. E-mail address: jmukho28@gmail.com (J. Mukhopadhyay).

https://doi.org/10.18231/j.jchm.2023.006 2394-272X/© 2023 Innovative Publication, All rights reserved.

2. Materials and Methods

The study was conducted by purposive selection of all COVID-19 affected individuals who came in contact with the incumbent through personal and professional threads of communication during the pandemic. 83 such individuals, who were declared COVID-19 positive by RT-PCR in Govt. & Private hospitals and clinics during May 20 to Sep 21, were contacted through e-mode for enlisting and online individual consent, however only 79 of them did respond. Sample size was considered on the basis of incidence of COVID-19 in Kolkata (2.26%) considering total cases in 2021 as documented 3.37 Lacs in a population of 14.9 Million approximating to a sample of 36 at 5% error.⁴ However, as 79 COVID affected subjects registered for the study, all were included. A structured self-admissible organised questionnaire was prepared after scrutinizing available literature and international scientific data on the issue which included the participants' personal attributes, details of COVID-19 infection and the post-COVID medico-social ordeal experienced by them. The equestion-naires were distributed e-mode/online sequentially as the incumbent came in contact with the affected subjects during the study period. The participants were contacted to note their conditions every week during first month, thereafter once in 4 weeks till 12 weeks. The filledin questionnaires received initially were topped-up with information received later during weekly/monthly follow up of each subject. Finally, questionnaires were scrutinized, analyzed and tabulated. Socio-economic status (SES) was ascertained as per current scale.⁵ Common statistical applications were used to determine significance as needed appropriate.

3. Results

There were 34 & 45 individuals who contracted COVID-19 in 2020 & 2021 respectively (Table 1). Majority of the subjects belonged to 41-50 years age group (47% & 44%) in both the groups followed by 31-40 years group (26.5%) & 26.7%). Large percentage were Hindu (91%) male (60% & above) and educated to post-graduate level (around 67%) in both the clusters. 79% & 93% were employed as stated; significantly large no. of affected (23.5%) lost job during 2020. 85-88% of the employed were benefitted by jobs in Govt. sector being benched with full pay. 18 (52.9%) & 16 (35.6%) respondents had co-morbidities respectively, of which hypertension in the first group and coronary artery disease (CAD) in the second were predominant. A little above 35% in each group contracted COVID-19 in the month of May'20 & '21 and Jun '21 respectively. Significantly higher proportions (88%) were admitted in hospital in first wave as compared to the second (69%). Mostly the contacts of both sets of patients were quarantined at home and significantly higher proportions (47%) of the first group had one or other addiction.

Cent-percent respondents of both the assemblages practiced washing hands twice daily before meals, wearing mask outdoor, social distancing, up-keeping personal hygiene, using sanitizer, observing cough/sneeze discipline and avoiding personal contact greetings. However, frequent hand washing (>3 times a day) & using gloves outdoor were not in that high order.

During the first wave, majority (100%) experienced fever, weakness, breathlessness and cough with sore throat during first 3 weeks, continued to experience the same (except fever) till 9 weeks (94-100%) and many of them (35-82%) remained symptomatic beyond 9 weeks following the initial acquisition of COVID-19 (Table 3). Headache (94%), anosmia & ageusia (64.7%), giddiness, weight-gain & sleeplessness (35% each) constituted other persisting symptoms that even continued in the proportion of 29 to 76% till 9 weeks later resolved to certain extent except for anosmia-ageusia, weight gain and insomnia. Diarrhoea, irritability, arthralgia (29% each) and mental clouding (23.5%) were among other important symptoms during first 3 weeks that persisted approximately in the same proportion till 8 weeks and faded away thereafter except for arthralgia & irritability continuing beyond 9 weeks.

During the second surge in 2021, majority (100%) experienced fever, weakness, breathlessness followed by cough with sore throat (93,3%) and headache (91%) during first 3 weeks, continued to experience the same except fever till 9 weeks (89-100%) and considerable proportion (40-89%) remained symptomatic beyond 9 weeks following COVID-19 infection. Anosmia & ageusia (60%), insomnia, diarrhoea & arthralgia (40% each) represented other lasting symptoms that even continued in the proportion of 13 to 48.8% till 9 weeks later got settled except for loss of smell/ taste and insomnia. Mood swing, weight gain (33% each) with giddiness & mental clouding (20% each) were relevant in the first 3 week, continually experienced by 20% of the affected subjects till 9 weeks in 13% patients.

During first wave, isolation, anxiety, avoidance by relatives, difficulties in procuring common items & medicines along with apathy from attendants/drivers became part of life for cent-percent affected subjects in the first 3 weeks after COVID infection that extended till 9 weeks for 76% & above respondents, further remained as reality for existence for 59-64.7% of the patients beyond 9 weeks (Table 4). Many (91%) felt lonely, disinterested in surroundings and lost-access to health care during first 3 weeks finally continuing among 29-59% of the subjects even beyond 9 weeks. 35% stated about financial constraints due to extra expenditure and feeling of helplessness till 9 weeks and 26.5% had the same ahead of 9 weeks also.

Personal Attributes	Sub-attributes	Contracted in 2020 (n1-34)	Contracted in 2021 (n2-45)	Significance
	20 - 30 years	3 (8.82)	5(11.11)	
Age	31 - 40 years	9(26.47)	12(26.67)	Chi 0 1206 NS
(n-79)	41 - 50 years	16(47.05)	20(44.44)	Cm-sq = 0.1296, NS
	51 years and above	6(17.65)	8(17.78)	
Gender	Male	21(61.76)	27(60.0)	Chi az = 0.0252
(n-79)	Female	13(38.24)	18(40.0)	Cm-sq = 0.0233
D-1:-:	Hindu	31(91.18)	41(91.11)	
(n, 70)	Muslim	3(8.82)	3(6.67)	Chi-sq – 0.0001, NS
(11-79)	Others	-	1(2.22)	
Education	Graduate	11(32.35)	15(33.33)	Chi ca = 0.008 NS
(n-79)	Post-graduate	23(67.65)	30(66.67)	Cm-sq = 0.008, NS
Working status (p. 70)	Employed	26(76.45)	42(93.33)	Chi-sq – 4.59,
working status (n-79)	Lost job	8(23.53)	3(6.67)	p<0.05
Working org (n-68)	Govt. services	22(84.61)	36(85.71)	Chi ca = 0.002 NS
(n1-26, n2-42)	Private org	4(15.38)	6(14.29)	Cm-sq = 0.003, NS
SES (n-79)	Higher	29(85.30)	38(84.44)	Chi ag 0.01 NS
	Upper middle	5(14.70)	7(15.56)	CIII-Sq = 0.01, NS
Co monhidition	Hypertension	9(50.00)	4(25.00)	
(n, 26)	Diabetes	7(38.89)	3(18.75)	Chi-sq – 7.89,
(n^{-50}) (n^{-1})	CAD	1(5.56)	5(31.25)	p<0.05
(11-10, 112-10)	Others	1(5.56)	4(25.00)	
	May	2(5.88)	16(35.56)	
Months of COVID	Jun	3(8.82)	16(35.56)	Chi az = 20.38
Acquisition	July	8(23.52)	7(15.56)	r = 29.38,
(n-79)	Aug	9(26.47)	5(11.11)	p<.0001
	Sep	12(35.29)	1(2.22)	
Admitted to Hospital (n-7	79)	30(88.24)	31(68.89)	Chi-sq – 4.12, p<0.05
Quarantine of contacts	Home	34(91.89)	48(96.0)	Chi = 0.66 MS
(n-87)	Center	3(8.11)	2(4.0)	Cn1-sq = 0.66, NS
$(n_{1,3,7}, n_{2}-50)$	Tobacco	7(20.59)	5(11.11)	
(n-79)	Alcohol	9(26.47)	6(13.33)	P1-47.1, P2-24.4, 7.2, 12, m < 0.05
(n1-34, n2- 45)	Others	-	-	Z-2.12, p<0.05

Table 1: Socio-demographic attributes of COVID-19 patient	ients

Table 2: Practice of COVID appropriate measures

COVID Appropriate Bahaviour	Followed precautions			
COVID Appropriate behaviour	2020 (n-34)	2021 (n-45)		
	No. (%)	No. (%)		
Washing hand with soap/water twice, at least before meals	34(100.0)	45(100.0)		
Washing hand with soap & water 3 times or more everyday	20(58.82)	24(53.33)		
Wearing mask while going out for daily need or work	34(100.0)	45(100.0)		
Social distancing from others in shops, markets & public	34(100.0)	45(100.0)		
places				
Wearing gloves in market places & shops	20(58.82)	24(53.33)		
Taking daily bath & maintaining personal hygiene	34(100.0)	45(100.0)		
Using alcohol based sanitizer to disinfect hand	34(100.0)	45(100.0)		
Covering mouth & nose in case of cough & sneezes	34(100.0)	45(100.0)		
Not shaking hand or hugging anyone	34(100.0)	45(100.0)		

	Con	Contracted in 2020 (n-34)			Contracted in 2021 (n-45)		
Medical Problems	No. of patients had symptoms (%)		No. of patients had symptoms (%)				
	03 weeks	04-09 weeks	> 09 weeks	03 weeks	04-09 weeks	> 09 weeks	
Moderate to high Fever	34 (100.00)	8 (23.52)	-	45 (100.00)	6(13.33)	-	
Weakness/tiredness	34 (100.00)	34 (100.00)	28 (82.35)	45 (100.00)	45 (100.00)	40 (88.88)	
Breathlessness	34 (100.00)	32 (94.11)	26 (76.47)	45 (100.00)	42 (93.33)	40 (88.88)	
Cough/sore throat	34 (100.00)	32 (94.11)	12 (35.29)	42 (93.33)	42 (93.33)	36 (80.00)	
Headache	32 (94.11)	26 (76.47)	-	41 (91.11)	40 (88.88)	18 (40.0)	
Loss of smell & Taste	22 (64.70)	20 (58.82)	10 (29.41)	27 (60.0)	22 (48.88)	6 (13.33)	
Giddiness	12 (35.29)	10 (29.41)	-	9 (20.0)	-	-	
Weight gain	12 (35.29)	10 (29.41)	10 (29.41)	15 (33.33)	9 (20.00)	-	
Sleep problem	12 (35.29)	10 (29.41)	9 (26.47)	18 (40.00)	9 (20.00)	9 (20.00)	
Diarrhoea	10 (29.41)	9 (26.47)	-	18 (40.00)	6 (13.33)	-	
Change in mood	10 (29.41)	10 (29.41)	10 (29.41)	15 (33.33)	9 (20.00)	6 (13.33)	
Arthralgia/Myalgia	10 (29.41)	10 (29.41)	8 (23.52)	18 (40.00)	18 (40.00)	6 (13.33)	
Mental clouding	8 (23.52)	8 (23.52)	-	9 (20.00)	9 (20.00)	6 (13.33)	

Table 3: Persistence of medical symptoms of COVID patients

Table 4: Social set-backs experienced by COVID patients

	Contracted in 2020 (n-34)		Contracted in 2021 (n-45)			
Social Problems	No.(%) expressed difficulties		No.(%) expressed difficulties			
	03 weeks	04-09 weeks	>09 weeks	03 week	04-09 weeks	>09 weeks
Isolation	34 (100.0)	26 (76.47)	22 (64.70)	45 (100.0)	42 (93.33)	36 (80%)
Anxiety for persistent symptoms	34 (100.0)	26 (76.47)	22 (64.70)	45 (100.0)	42 (93.33)	36 (80%)
Avoidance by relatives & Friends	34 (100.0)	26 (76.47)	22 (64.70)	45 (100.0)	42 (93.33)	36 (80%)
Difficulty in procuring common medicines	34 (100.0)	26 (76.47)	22 (64.70)	45 (100.0)	42 (93.33)	36 (80%)
Avoidance by attendants, drivers & domestic help	34 (100.0)	26 (76.47)	20 (58.82)	42 (93.33)	40 (88.88)	27 (60.0)
Difficulty in going out for provisions	34 (100.0)	28 (82.35)	20 (58.82)	42 (93.33)	40 (88.88)	27 (60.0)
Loneliness	31(91.17)	24 (70.59)	20 (58.82)	42 (93.33)	40 (88.88)	27 (60.0)
Loss of interest in Surrounding	31(91.17)	24 (70.59)	20 (58.82)	40 (88.88)	36 (80%)	27 (60.0)
Loss of access to health care	31(91.17)	24 (70.59)	10 (29.41)	40 (88.88)	36 (80%)	22 (48.88)
Financial constraint	12 (35.29)	12 (35.29)	9 (26.47)	9 (20.0)	9 (20.0)	6 (13.33)
Helplessness	12 (35.29)	12 (35.29)	9 (26.47)	9 (20.0)	9 (20.0)	-

During second surge, cent-percent stated social barriers of isolation, anxiety, avoidance by relatives and trouble in obtaining medicines in the first 3 weeks after COVID acquisition that persisted till 9 weeks for 93% of the afflicted, remained further among 80% of the patients beyond 9 weeks. 93% exclaimed of loneliness, avoidance by attendants/driver and difficulty in getting daily provisions during first 3 weeks even suffering through the same hurdle in high proportion (89%) till 9 weeks, scaled down to 60% past 9 weeks after COVID infection. 89% revealed loss of interest in surrounding and difficulty in accessing medical care during first 3 weeks truncated to 80% & 49% sequentially later till 9 weeks and beyond. 20% reiterated about financial constraints and helplessness all through 9 weeks after initial infection.

All respondents during both the phases of pandemic disclosed the wish of spending more time with family, abatement of anxiety, requirement of a counselor and

Perceived support/need for social problems	Contracted in 2020 (n-34) No. (%)	Contracted in 2021 (n-45) No. (%)
More time needed with working spouse at home	34(100.0)	45(100.0)
Someone who could absolve anxiety	34(100.0)	45(100.0)
Requirement of a counselor/professional	34(100.0)	45(100.0)
Online counselling support	32(94.11)	45(100.0)
Streamlining of domestic support	34(100.0)	42(93.33)
Reformation of home delivery & procurement	34(100.0)	45(100.0)
Relaxation in containment rule & dispel stigma	31(91.17)	40(88.88)

Table 5: Perceived need for social support

restructuring of home delivery system. Large contingent during the first wave insisted online counseling assistance (94%) and cutback in containment rules (91%) to dispel stigma as against 89 to 93% of second surge, demanding domestic support for elderly and curtailment of containment rules.

4. Discussion

The study included patients from both first and second surge of COVID-19 pandemic in Kolkata, India during 2020 & 2021. Affected individuals belonged to different age groups, educational and economic status. Majority belonged to 41-50 years age group (47% & 44% respectively) followed by 31-40 years group (above 26% each) in both the phases of pandemic. Preponderantly educated employed males (60% & above) were affected indicating their possible vulnerability to exposure to infectious environment. A recent study documented consistent predominance of males (65.39%) compared to females (34.61%) among COVID-19 patients in all age categories.⁶ The researchers consider higher rates of smoking, lesser hand washing, existing respiratory illness and biological difference between sexes as driving force for higher infection rate among males.⁷ Preventive measures to avoid COVID-19 were found significantly associated with the gender where females noticed to have been more strict followers compared to males.8

Co-morbidities existed in 52.9% & 35.6% respondents respectively of the corresponding phases, of which hypertension in the first group and CAD in the second found the leading. Recent studies documented that 20-51%of patients of COVID-19 reportedly having one or more comorbidity with diabetes (10–20%), hypertension (10–15%), cardio and cerebro-vascular diseases (7–40%) being the common.^{9,10}

A little above 35% in each phase got COVID-19 infection in the month of Sep'20 & May and Jun '21 respectively. Reportedly crests of epidemicity were reached in these months in India during the year'20 & '21 respectively.¹¹ Seasonal variations has been well-established criteria for many respiratory viral infections including COVID-19 indicating association between temperature, humidity, UV radiations and contagion presence; however other analyses have failed to show a credible role of climate and weather particularly when population immunity is low.^{12,13} Governmental non-pharmaceutical interventions (NPIs) like lock-down, closure of Govt. offices, educational institutions, malls & movie halls resulted in waxing and waning of infection rate in consecutive waves since early 2020, effect of which could be crucial than environmental determinants.¹⁴ Significantly 88% of the subjects were admitted in hospital during first wave as compared to the second (69%) however, there were no fatalities. During the second wave in India, the proportion of active cases that needed hospital care remained generally in the range of 20-23 percent.¹⁵ Most contacts of both groups were quarantined at home and significantly 47% patients of the first phase had one or other addictions. Socio-demographic traits and the dynamics of the disease transmission among the subjects in the present endeavour showed commonality in most of the facets with recently published works probably asserting convincing internal validity of the sample.

Entire respondents of both the lots practiced washing hands twice daily before meals, wearing mask outdoor, social distancing, maintaining personal hygiene, using sanitizer, observing cough/sneeze discipline and avoiding contact greetings. However, frequent hand washing (>3 times a day) and use of gloves outdoor were not followed in that high order in the groups.

Authors ascribed around 95-98% of the educated subjects displayed correct preventive knowledge to avert COVID-19 following personal protective measures (PPM).¹⁶ It has been annotated that large no. of subjects followed PPM adequately by using masks (97.2%), social distancing (95.3%) and hand hygiene (91.2%) that apparently corroborates the observations of present work.¹⁵ Strong governmental initiatives coupled with synergistic effects of awareness, acceptance and action by the people made it possible to execute COVID appropriate behaviour

(CAB) satisfactorily to counter the spikes of COVID-19 infection. $^{\rm 17}$

During the first wave, everyone had fever, fatigue, breathlessness, cough & sore throat during first 3 weeks, continued (except fever) till 9 weeks (94-100%) and many (35-82%) remained symptomatic beyond 9 weeks following the infection. Headache (94%), anosmia & ageusia (64.7%), giddiness, weight-gain & sleeplessness (35% each) were persistent & lingered in high order (29-76%) till 9 weeks; further stuck around except for headache. Diarrhoea, irritability, arthralgia (29% each) and mental clouding (23.5%) were significant during first 3 weeks, persisted in similar order till 8 weeks and faded away subsequently except for arthralgia & irritability continuing beyond 9 weeks. Similar sort of manifestations were experienced for fairly same duration even in the second surge.

Reportedly at 4 months after initial infection, shortness of breath (8.6%), anosmia (12.4%), ageusia (11.1%), and fatigue (9.7%) persisted among the COVID-19 patients; generally one characteristic symptom was present in 27.8 & 34.8% at 4 and 7 months post-infection respectively.¹⁸ Anosmia and diarrhoea during acute COVID-19 were associated with higher risk of developing long-term symptoms. In line with data published by Townsend et al, it was seen that even patients with initially mild disease may develop fatigue as a leading and debilitating feature making individual non-participating in everyday & work-life.¹⁹

Moreno-Perez et al reported that up to 50.9% of the patients considered "recovered" even with persisted symptoms similar to those experienced during the acute phase.²⁰ Moreover, other studies showed that the reported duration of myalgia-arthralgia varies widely depending on duration of patients' follow-up.^{21–23}

Isolation, anxiety, avoidance by relatives, hardships in procuring daily needs & medicines along with indifference from attendants/drivers were of high order (93-100%) among the indisposed of both phases of pandemic during first 3 weeks existing in the range of 59-80% even beyond 9 weeks after COVID acquisition. Loneliness, loss of interest in the daily errands and lost-access to health care during first 3 weeks were considerably high (89-91%) further extending in the range of 29-60% past 9 weeks among the affected of both the upsurges. Financial constraints due to added expenses and helplessness were experienced in the ambit of 13-35% among the patients of both stages of catastrophe over entire period.

Studies reported that patients with COVID-19 had symptoms of anxiety (42%), depression (31%) and insomnia (40%).^{24,25} Recent work disclosed that post-traumatic stress associated with the COVID-19 was 96.2% among COVID-19 patients and half of the participants considered psycho-educational services helpful.²⁶ Zhang et al documented that depression (29.2%) was quite predominant in COVID-19 patients associated with anxiety-

driven behavior like easy annoyance & irritability.²⁷ Findings of the present intent on social setbacks is higher than the documented observations and that appears to be due to the fact that all these factors have been self-stated subjective reflection of affected psyche with immediate recollection resulting in higher cogitation; therefore needs guarded consideration while comparing with findings of other studies.

The study revealed that more time with family, alleviation of anxiety through counselor and revamping of home delivery services were of high demand (100%) together with online support and relaxation in containment rules (>89%) among the affected of both phases. The preventive measures of isolation, quarantine and containment related restricted mobility for the patients & their contacts, adopted in managing the pandemic showed unwanted consequences like anxiety, depressive episodes and post-traumatic stress among the sensitive subjects; when prolonged, did aggravate negative emotions and psychological breakdown.²⁸ Promotion of psychological interventions on the specific population who is more likely to develop negative emotions and suffering is needed. The Lancet Global Mental Health Commission reported that the use of digital technologies can provide mental health interventions in order to reduce anxiety & stress and increase self-efficacy.^{29,30} It has been proclaimed that online counseling bypasses geographical barriers and is suitable to become a useful integrated tool in addition to traditional psychotherapy.³¹ Solidarity & compassion including food & medicine supply for isolated afflicted by generating a feeling of 'forced empathy' need to be accentuated by the use of hi-tech gadgets & means that may assist in depersonalization of associations & relationships, forcing the sense of closeness & familiarity, at least virtually. The hyper-connection of feelings & emotional bonding becomes a way to dissipate isolation and its consequences.

5. Conclusion and Limitation

This has been a controlled work carried out among the linked patients of the incumbent scholar, who were generally contacted through telecom & e-modes during strict surveillance with restricted mobility phase of the pandemic in Kolkata. Therefore it needs careful consideration for generalization and comparison with other literature. However, the socio-demographic traits of the affected including disease transmissibility patterns were similar to recently documented literatures, possibly pointing to a plausible coherent sample. The study unveiled prolonged sufferings of the COVID-19 affected in medicalsocial-diurnal front especially among the elders and directs activation of online services for medical, social, psychological & humanitarian support & assistance. Further research in comparable directions may lend a hand to decide prospective line of approach to resolve the tribulations of epidemic-affected people in future comprehensively.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

- Chopra V, Flanders SA, O'Malley M, Malani AN, Prescott HC. Sixtyday outcomes among patients hospitalized with COVID-19. *Ann Intern Med.* 2020;174(4):576–8.
- Goertz YMJ, Herck MV, Delbressine JM, Vaes AW, Meys R, Machado FVC, et al. Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome? *ERJ Open Res.* 2020;6(4):00542–2020.
- Mandal S, Barnett J, Brill SE, Brown JS, Denneny EK, Hare SS, et al. Long-COVID: A cross-sectional study of persisting symptoms, biomarker and imaging abnormalities following hospitalisation for COVID-19. *Thorax.* 2021;76(4):396–8.
- West Bengal COVID-19 Health Bulletin; Dec 2021. Available from: https://www.wbhealth.gov.in/uploaded_files/corona/WB_DHFW_ Bulletin_31ST_DEC_REPORT_FINAL.pdf.
- Pandey VK, Aggarwal P, Kakkar R. Modified BG Prasad's Socioeconomic Classification - 2018: The need of an update in the present scenario. *Indian J Community Health*. 2018;30(1):82–4.
- Kushwaha S, Khanna P, Rajagopal V, Kiran T. Biological attributes of age and gender variations in Indian COVID-19 cases: A retrospective data analysis. *Clin Epidemiol Glob Health*. 2021;11:100788.
- Betron M, Gottert A, Pulerwitz J, Shattuck D, Stevanovic-Fenn N. Men and COVID-19: adding a gender lens. *Glob Public Health*. 2020;15(7):1090–2.
- Tegegne KT, Assefa AA, Tegegne ET, Tessema MK. Implication of gender on Covid-19 prevention practice among adults in Ethiopia: A systemic review and meta-analysis. *Res Rev Health Care OAJ*. 2020;6(1):562–6.
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y. Clinical features of patients with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497–506.
- Chen N, Zhou M, Dong X. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395:507–513.
- 11. World-Countries-India. 2022.
- Carlson CJ, Gomez ACR, Bansal S, Ryan SJ. Misconceptions about weather and seasonality must not misguide COVID-19 response. *Nat Commun.* 2020;11(1):1–4. doi:10.1038/s41467-020-18150-z.
- Baker RE, Yang W, Vecchi GA, Metcalf CJE, Grenfell BT. Susceptible supply limits the role of climate in the early SARS-CoV-2 pandemic. *Science*. 2020;369(6501):315–9.
- Mcclymont H, Hu W. Weather Variability and COVID-19 Transmission: A Review of Recent Research. *Int J Environ Res Public Health.* 2021;18(2):396.
- The Indian Express. Covid hospitalisation rate at 5-10% but situation may change soon: Govt; 2022. Available from: https://indianexpress.com/article/india/covid-omicronhospitalisation-rate-low-govt-7716156/.
- Shukla S, Deotale P. Knowledge, attitude and practices towards COVID-19 pandemic in the community: a cross-sectional web-based survey in India. *Int J Res Med Sci.* 2020;8(10):3652–56.

- Tomar BS, Singh P, Suman S, Raj P, Nathiya D. Indian community's knowledge, attitude and practice towards COVID-19. *MedRxiv*. 2020;p. 1–20. doi:10.1101/2020.05.05.20092122.
- Augustin M, Schommers P, Stecher M, Dewald F, Gieselmann L, Gruell H, et al. Post- COVID syndrome in non-hospitalised patients with COVID-19: A longitudinal prospective cohort study. *Lancet Reg Health Eur.* 2021;6:100122.
- Townsend L, Dyer AH, Jones K, Dunne J, Mooney A, Gaffney F, et al. Persistent fatigue following SARS-CoV-2 infection is common and independent of severity of initial infection. *PLoS ONE*. 2020;15(11):e0240784.
- Moreno-Pérez O, Merino E, Leon-Ramirez JM, Andres M, Ramos JM, Arenas-Jiménez J, et al. Post-acute COVID-19 syndrome. Incidence and risk factors: A Mediterranean cohort study. *J Infect*. 2021;82(3):378–83.
- Arnold DT, Hamilton FW, Milne A, Morley AJ, Viner J, Attwood M, et al. Patient outcomes after hospitalisation with COVID-19 and implications for follow-up: results from a prospective UK cohort. *Thorax*. 2020;76(4):399–401.
- Carfi A, Bernabei R, Landi F. Persistent symptoms in patients after acute COVID- 19. JAMA. 2020;324(6):603–5.
- Jacobs LG, Paleoudis EG, Bari DLD, Nyirenda T, Friedman T, Gupta A, et al. Persistence of symptoms and quality of life at 35 days after hospitalization for COVID-19 infection. *PLoS One*. 2020;15(12):e0243882.
- Paz C, Mascialino G, Adana-Díaz L, Rodríguez-Lorenzana A, Simbaña-Rivera K, Gómez-Barreno L, et al. Anxiety and depression in patients with confirmed and suspected COVID-19 in Ecuador. *Psychiatry Clin Neurosci*. 2020;74(10):554–5.
- Mazza GM, DeLorenzo R, Conte C, Poletti S, Vai B, Bollettini I, et al. Anxiety and depression in COVID-19 survivors: role of inflammatory and clinical predictors. *Brain Behav Immun.* 2020;89:594–600.
- Bo HX, Li W, Yang Y, Wang Y, Zhang Q, Cheung T, et al. Posttraumatic stress symptoms and attitude toward crisis mental health services among clinically stable patients with COVID-19 in China. *Psychol Med.* 2021;51(6):1052–3.
- Zhang J, Lu H, Zeng H, Zhang S, Du Q, Jiang T, et al. The differential psychological distress of populations affected by the COVID-19 pandemic. *Brain Behav Immun.* 2020;87:49–50.
- Saladino V, Algeri D, Auriemma V. The Psychological and social impact of Covid-19: New perspectives of well-being. *Front Psychol.* 2020;11:577684. doi:10.3389/fpsyg.2020.577684.
- Patel V, Saxena S, Lund C, Thornicroft SG, Baingana F, Bolton P, et al. The Lancet Commission on global mental health and sustainable development. *Lancet J*. 2018;392(10157):1553–98.
- Xiao H, Zhang Y, Kong D, Li S, Yang N. The effects of social support on sleep quality of medical staff treating patients with coronavirus disease (COVID-19) in January and February 2020 in China. *Med Sci Monit*. 2020;26:e923549.
- Perrin P, Rybarczyk B, Pierce B, Jones H, Shaffer C, Islam L. Rapid tele-psychology deployment during COVID-19 pandemic: A special issue commentary and lessons from primary care psychology training. *J Clin Psychol*. 2020;76(6):1173–85.

Author biography

J Mukhopadhyay, Professor (1) https://orcid.org/0000-0002-9470-9453

Cite this article: Mukhopadhyay J. Post-COVID impact: Misery of the indisposed. *J Community Health Manag* 2023;10(1):25-31.